



Government Polytechnic, Ahmedabad

Science and Humanities Department

Engineering Mathematics

Subject Code: 4320002

Unit -03 Integration and Its Applications

[Marks – 14]

Course Outcome (CO c):

Demonstrate the ability to solve engineering related problems based on applications of integration.

Important formula of Integration

$$(1) \int x^n dx = \frac{x^{n+1}}{n+1} + c ; n \neq 1$$

$$(2) \int 1 dx = x + c$$

$$(3) \int \frac{1}{x} dx = \log_e x + c$$

$$(4) \int k dx = kx + c ; \text{ where } k \text{ is constant}$$

$$(5) \int e^x dx = e^x + c$$

$$(6) \int a^x dx = \frac{a^x}{\log_e a} + c$$

$$(7) \int \sin x dx = -\cos x + c$$

$$(8) \int \cos x dx = \sin x + c$$

$$(9) \int \tan x dx = \log|\sec x| + c$$

$$(10) \int \cot x dx = \log|\sin x| + c$$

$$(11) \int \csc x dx = \log|\csc x - \cot x| + c$$

$$= \log \left| \tan \frac{x}{2} \right| + c$$

$$(12) \int \sec x dx = \log|\sec x + \tan x| + c$$

$$= \log \left| \tan \left(\frac{\pi}{4} + \frac{x}{2} \right) \right| + c$$

$$(13) \int \sec^2 x \, dx = \tan x + c$$

$$(14) \int \csc^2 x \, dx = -\cot x + c$$

$$(15) \int \sec x \cdot \tan x \, dx = \sec x + c$$

$$(16) \int \csc x \cot x \, dx = -\csc x + c$$

$$(17) \int \frac{1}{\sqrt{a^2 - x^2}} \, dx = \sin^{-1} x + c \quad or \quad -\cos^{-1} x + c$$

$$(18) \int \frac{1}{\sqrt{1-x^2}} \, dx = \sin^{-1} x + c \quad or \quad -\cos^{-1} x + c$$

$$(19) \int \frac{1}{x^2+a^2} \, dx = \frac{1}{a} \tan^{-1} \frac{x}{a} + c$$

$$(20) \int \frac{1}{1+x^2} \, dx = \tan^{-1} x + c \quad or \quad -\cot^{-1} x + c$$

$$(21) \int \frac{1}{|x|\sqrt{x^2-a^2}} \, dx = \frac{1}{a} \sec^{-1} \frac{x}{a} + c \quad or \quad \frac{-1}{a} \csc^{-1} \frac{x}{a} + c$$

$$(22) \int \frac{1}{x\sqrt{x^2-1}} \, dx = \sec^{-1} x + c \quad or \quad -\csc^{-1} x + c$$

$$(23) \int \frac{1}{\sqrt{x^2+a^2}} \, dx = \log \left| x + \sqrt{x^2+a^2} \right|$$

$$(24) \int \frac{1}{\sqrt{x^2-a^2}} \, dx = \log \left| x + \sqrt{x^2-a^2} \right|$$

$$(25) \int \frac{1}{x^2-a^2} \, dx = \frac{1}{2a} \log \left| \frac{x-a}{x+a} \right| + c \quad ; \quad (x^2 > a^2)$$

$$(26) \int \frac{1}{a^2-x^2} \, dx = \frac{1}{2a} \log \left| \frac{a-x}{a+x} \right| + c \quad ; \quad (x^2 < a^2)$$

$$(27) \int [f(x)]^n f'(x) dx = \frac{[f(x)]^{n+1}}{n+1} + c ; n \neq -1$$

$$(28) \int \frac{f'(x)}{f(x)} dx = \log|f(x)| + c$$

Working rules of Integration

$$(29) \int [f(x) + g(x)] dx = \int f(x) dx + \int g(x) dx$$

$$(30) \int [f(x) - g(x)] dx = \int f(x) dx - \int g(x) dx$$

$$(31) \int k.f(x) dx = k \int f(x) dx + c; k \text{ is constant}$$

Integration by Parts

$$(32) \int u \cdot v dx = u \int v dx - \int \left(\frac{du}{dx} \int v dx \right) dx + c$$

$$(33) \int \sqrt{x^2 + a^2} dx = \frac{x}{2} \sqrt{x^2 + a^2} + \frac{a^2}{2} \log |x + \sqrt{x^2 - a^2}| + c$$

$$(34) \int \sqrt{x^2 - a^2} dx = \frac{x}{2} \sqrt{x^2 - a^2} + \frac{a^2}{2} \log |x + \sqrt{x^2 - a^2}| + c$$

$$(35) \int \sqrt{a^2 - x^2} dx = \frac{x}{2} \sqrt{a^2 - x^2} + \frac{a^2}{2} \sin^{-1} \frac{x}{a} + c ; (a > 0)$$

$$(36) \int e^x [f(x) + f'(x)] dx = e^x f(x) + c$$

$$(37) \int e^{ax} \sin bx \, dx = \frac{e^{ax}}{a^2 + b^2} (a \sin bx - b \cos bx) + c$$

$$(37) \int e^{ax} \cos bx \, dx = \frac{e^{ax}}{a^2 + b^2} (a \cos bx + b \sin bx) + c$$

Important Trigonometric Substitution

No.	Integrand	Substitution
1	$a^2 - x^2$ <u>or</u> $\sqrt{a^2 - x^2}$	$x = a \sin \theta \quad or \quad x = a \cos \theta$
2	$x^2 - a^2$ <u>or</u> $\sqrt{x^2 - a^2}$	$x = a \sec \theta \quad or \quad x = a \cosec \theta$
3	$x^2 + a^2$ <u>or</u> $\sqrt{x^2 + a^2}$	$x = a \tan \theta \quad or \quad x = a \cot \theta$
4	$\sqrt{a + x}$	$x = a \tan^2 \theta \quad or \quad x = a \cos 2\theta$
5	$\sqrt{a - x}$	$x = a \sin^2 \theta \quad or \quad x = a \cos 2\theta$
6	$\sqrt{\frac{a - x}{a + x}}$	$x = a \cos 2\theta$
7	$\sqrt{2ax - x^2}$	$x = 2a \sin^2 \theta$

Important Trigonometric Formula

Identity No. 1: $\sin^2 \theta + \cos^2 \theta = 1$

$$\rightarrow \sin^2 \theta = 1 - \cos^2 \theta$$

$$\rightarrow \cos^2 \theta = 1 - \sin^2 \theta$$

Identity No. 2: $\tan^2 \theta + 1 = \sec^2 \theta$ (We get it by dividing $\cos^2 \theta$ to Identity No.1)

$$\rightarrow \tan^2 \theta = \sec^2 \theta - 1$$

$$\rightarrow \tan^2 \theta - \sec^2 \theta = 1$$

Identity No. 3: $1 + \cot^2 \theta = \operatorname{cosec}^2 \theta$ (We get it by dividing $\sin^2 \theta$ to Identity No.1)

$$\rightarrow \cot^2 \theta = \operatorname{cosec}^2 \theta - 1$$

$$\rightarrow \operatorname{cosec}^2 \theta - \cot^2 \theta = 1$$

Properties of Definite Integration

Even Function:

If $f(-x) = f(x)$; for all x

e.g. (1) $\cos x$, (2) $\sec x$

(3) x^2 (4) x^4 (5) $x^{(\text{even number})}$

Odd Function:

If $f(-x) = -f(x)$; for all x

e.g. (1) $\sin x$, (2) $\tan x$ (3) $\operatorname{cosec} x$, (4) $\cot x$

(5) x (6) x^3 (7) $x^{(\text{odd number})}$

Even Function \times *Even Function* = *Even Function*

Odd Function \times *Even Function* = *Odd Function*

Even Function \times *Odd Function* = *Odd Function*

Odd Function \times *Odd Function* = *Even Function*

e.g.

$x^2 \cos x$ is an *even* function

$x \cos x$ is an *odd* function

$x^2 \sin x$ is an *odd* function

$x \sin x$ is an *even* function

$$\int_{-a}^a f(x) dx = \begin{cases} 2 \int_0^a f(x) dx; & \text{if } f(x) \text{ is an even function} \\ 0; & \text{if } f(x) \text{ is an odd function} \end{cases}$$

$$\int_a^b f(x) dx = \int_0^b f(a+b-x) dx$$

$$\int_0^a f(x) dx = \int_0^a f(a-x) dx; \quad (\text{In above formula if } a=0 \text{ and } b=a)$$

$$\int_0^{2a} f(x) dx = \int_0^a f(x) dx + \int_0^a f(2a-x) dx;$$

Dr. Dushyant G. Patel

Assignment - 3

वी अग्रिम वार्षिक प्रश्नपत्र
Date: / / Pg.no.:

Unit - 03

Integration and its Applications:

* Question Set for 01 Marks:

$$1. \int 6x^5 dx = x^6 + C$$

$$\begin{aligned} & \int 6x^5 \\ & 6 \int x^5 dx \\ & \frac{6x^6}{6} \end{aligned}$$

$$\frac{x^6}{6}$$

$$x^6$$

$$(2) \int (\cos^2 x + \sin^2 x) dx = x + C$$

$$\rightarrow a) x$$

$$(\because \cos^2 x + \sin^2 x = 1)$$

$$(\because \int 1 dx = x + C)$$

$$(3) \int \frac{\cos x}{\sin x} dx = -\operatorname{cosec}^2 x + C$$

$$\rightarrow -\operatorname{cosec}^2 x$$

$$(4) \int a^x dx = \frac{a^x}{\log_e a} + C$$

$$\rightarrow \frac{a^x}{\log_e a} + C$$

5) $\int \frac{1}{1+x^2} dx = \tan^{-1} x + C$

\rightarrow c) $\tan^{-1} x$

6) $\int \frac{\log x}{x} dx = \frac{1}{2}(\log x)^2$

$\rightarrow \frac{1}{2}(\log x)^2$

($\because \log x = t \quad \int t \cdot dt$)

$\frac{1}{x} = \frac{dt}{dx} = \frac{t^2}{2} + C$

$\frac{dx}{x} = dt$

$\boxed{\int \frac{1}{2}(\log x)^2 + C}$

7) $\int_2^5 x^2 dx = 39 + C$

\rightarrow 9) 39

$\int_2^5 x^2 dx = \left[\frac{x^3}{3} \right]_2^5$

$= \frac{(5)^3 - (2)^3}{3}$

$= \frac{125}{3} - \frac{8}{3}$

$= \frac{125-8}{3}$

$= \frac{117}{3} = \boxed{39}$

8) $\int_0^1 \frac{2}{1+x^2} dx = \frac{\pi}{2} + C$

$\underline{\underline{=}} \quad 2 \int \left[\frac{1}{1+x^2} \right]_0^1$

$2 [\tan^{-1} x]_0^1$

$2 [\tan^{-1} 1 - \tan^{-1} 0]$

$2 (\frac{\pi}{4} - 0)$

$\boxed{\frac{\pi}{2}}$

$$\begin{aligned}
 \textcircled{9} \quad & \int_0^{\frac{\pi}{4}} \frac{1}{4+x^2} dx = \frac{\pi}{8} + C \quad \xrightarrow{\text{Soln}} \int_0^{\frac{\pi}{4}} \frac{1}{x^2+2^2} dx \\
 \rightarrow & \frac{\pi}{8} \quad = \left[\frac{1}{2} \tan^{-1}\left(\frac{x}{2}\right) \right]_0^{\frac{\pi}{4}} \\
 & = \frac{1}{2} \cdot \left[\tan^{-1}\frac{\pi}{8} - \tan^{-1}0 \right] \\
 & = \frac{1}{2} \left[\frac{\pi}{8} - 0 \right] \\
 & = \boxed{\frac{\pi}{16}}
 \end{aligned}$$

$$\begin{aligned}
 \textcircled{10}) \quad & \int_{-5}^5 x^3 dx = 0 + C \quad \xrightarrow{\text{Soln}} \left[\frac{x^4}{4} \right]_{-5}^5 \\
 \rightarrow & \textcircled{6} 0 \quad = (5)^4 - (-5)^4 \\
 & = \frac{625}{4} - \left(-\frac{625}{4} \right) \\
 & = \frac{0}{4} = 0
 \end{aligned}$$

* Question Set for 03 marks:

$$\begin{aligned}
 \textcircled{1}) \quad & \int \frac{x^2 + 5x + 6}{x^2 + 2x} dx \\
 & \xrightarrow{\text{Soln}} \int \frac{(x+3)(x+2)}{x(x+2)} dx \\
 & = \int \frac{x+3}{x} dx
 \end{aligned}$$

$$\begin{aligned}
 & \int \frac{x}{x} + \frac{3}{x} dx \\
 & = \int 1 dx + 3 \int \frac{1}{x} dx \\
 & = \boxed{x + 3 \ln|x| + C \text{ A.W.}}
 \end{aligned}$$

$$③ \int (4x^3 - \frac{1}{x} + \sin x - e^x) dx$$

$$\int (4x^3 - \frac{1}{x} + \sin x - e^x) dx$$

$$\int 4x^3 dx - \int \frac{1}{x} dx + \int \sin x dx - \int e^x dx$$

$$= 4x \cdot \frac{x^4}{4} - \log |x| + (-\cos x) - e^x$$

$$= [x^4 - \log |x| - \cos x - e^x + C] \quad \underline{\text{Ans}}$$

$$④ \int x \cdot (2x^2 + 3)^8 dx$$

$$\text{Let } \frac{d}{dx} 2x^2 + 3 = u$$

$$\frac{d}{dx} 2(2x) + 0 = \frac{dy}{dx}$$

$$2x dx = \frac{dy}{4}$$

$$\text{Now } \int x (2x^2 + 3)^8 dx = \frac{u^9}{36} + C$$

$$= \int u^8 \cdot \frac{dy}{4} = \frac{(2x^2 + 3)^9}{36} + C$$

$$= \frac{1}{4} \int u^8 dy$$

$$= \frac{1}{4} \int u^8 dy$$

$$= \frac{1}{4} \cdot \frac{u^9}{9} + C$$

Ans

$$(4) \int \frac{(1+x)e^x}{\sin^2(xe^x)}$$

Soln

$$\text{Let } xe^x = u$$

$$x \cdot \frac{d}{dx} e^x + e^x \frac{d}{dx} x = \frac{dy}{dx}$$

$$= x \cdot e^x + e^x(1) = \frac{dy}{dx}$$

$$= xe^x + e^x = \frac{dy}{dx}$$

$$= e^x(1+x) dx = du$$

Now

$$\int \frac{dy}{\sin^2 u}$$

$$= \int \frac{1}{\sin^2 u} du$$

$$= \int \csc^2 u + c$$

$$= -\cot u + c$$

$$= -\cot(xe^x) + c$$

$$= \boxed{-\cot(xe^x) + c} \quad \underline{\text{Ans}}$$

Q-5

$$\int e^{\tan x} \sec^2 x dx$$

Soln

$$\text{Let, } \tan x = u$$

$$\sec^2 x dx = du$$

Now

$$\int e^u \cdot du$$

$$= e^u + c$$

$$= \boxed{e^{\tan x} + c} \quad \underline{\text{Ans}}$$

$$\int \frac{3x+2}{2x^2+x+1} dx$$

$$\int \frac{3x+2}{2x^2+x+1} dx$$

$$= \int \frac{\frac{3}{4}(4x+1) + \frac{5}{4}}{2x^2+x+1} dx$$

$$\left\{ \int \frac{P'(x)}{F(x)} dx = \log |F(x)| + C \right.$$

$$= \frac{3}{4} \int \frac{4x+1}{2x^2+x+1} dx + \frac{5}{4} \int \frac{1}{2x^2+x+1} dx$$

$$= \frac{3}{4} \log |2x^2+x+1| + \frac{5}{4} \int \frac{1}{2(x^2+x/2+1/2)} du$$

$$= \frac{3}{4} \log |2x^2+x+1| + \frac{5}{8} \int \frac{1}{x^2+x/2+1/2} dx$$

$$= \frac{3}{4} \log |2x^2+x+1| + \frac{5}{8} \int \frac{1}{(x+1/4)^2 - 1/16 + 1/2} dx$$

$$= \frac{3}{4} \log |2x^2+x+1| + \frac{5}{8} \int \frac{1}{(x+1/4)^2 + 2/16} dx$$

$$\text{Let } x+1/4 = t$$

$$= \frac{3}{4} \log |2x^2+x+1| + \frac{5}{8} \int \frac{1}{(\sqrt{2}/4)^2 + t^2} dt$$

$$= \frac{3}{4} \log |2x^2 + x + 1| + \frac{5}{8} \cdot \frac{1}{\sqrt{7}/4} \tan^{-1} \left(\frac{x}{\sqrt{7}/4} \right)$$

$$\int \frac{1}{x^2 + a^2} dx = \frac{1}{a} \tan^{-1} \left(\frac{x}{a} \right) + C$$

$$= \frac{3}{4} \log |2x^2 + x + 1| + \frac{5}{8} \cdot \frac{4}{\sqrt{7}} \tan^{-1} \left(\frac{x+1/4}{\sqrt{7}/4} \right)$$

$$= \frac{3}{4} \log |2x^2 + x + 1| + \frac{5}{2\sqrt{7}} \tan^{-1} \left(\frac{4x+1}{\sqrt{7}} \right)$$

$$= \boxed{\frac{3}{4} \log |2x^2 + x + 1| + \frac{5}{2\sqrt{7}} \tan^{-1} \left(\frac{4x+1}{\sqrt{7}} \right) + C}$$

Q-7 $\int x \cdot e^{3x} dx$

Ans $\{ \int u \cdot v dx = u \cdot \int v dx - \int \left[\frac{du}{dx} \cdot \int v dx \right] dx \}$

Let $u = x, v = e^{3x}$

$$= x \cdot \int e^{3x} dx - \int \left[1 \cdot \int e^{3x} \right] dx$$

$$= x \cdot \frac{e^{3x}}{3} - \int \left[1 \cdot \frac{e^{3x}}{3} \right] dx$$

$$= \frac{xe^{3x}}{3} - \frac{1}{3} \int e^{3x} dx$$

મનુને પ્રાર્થના એ શક્તિશાળી હૃથીયાર છે.

$$= xe^{\frac{3x}{3}} - \frac{1}{3} e^{\frac{3x}{3}}$$

$$= \left[\frac{1}{3} xe^{3x} - \frac{1}{9} e^{3x} + c \right] \quad \underline{\text{Ans}}$$

Q-8 $\int x \tan^{-1} x \, dx$

$$\text{Let } u = \tan^{-1} x, v = x$$

$$\int u \cdot v \, dx = u \cdot \int v \, dx - \int \left[\frac{du}{dx} \cdot \int v \, dx \right] \, dx$$

$$= \tan^{-1} x \int x \, dx - \int \left[\frac{d}{dx} \tan^{-1} x \int x \, dx \right] \, dx$$

$$= \frac{x^2}{2} \tan^{-1} x - \int \left[\frac{1}{1+x^2} \cdot \frac{x^2}{2} \right] \, dx$$

$$= \frac{x^2}{2} \tan^{-1} x - \frac{1}{2} \int \frac{x^2}{1+x^2} \, dx$$

$$= \frac{x^2}{2} \tan^{-1} x - \frac{1}{2} \int \frac{(1+x^2)-1}{(1+x^2)} \, dx$$

$$= \frac{x^2}{2} \tan^{-1} x - \frac{1}{2} \left[\int 1 - \frac{1}{1+x^2} \, dx \right]$$

$$= \frac{x^2}{2} \tan^{-1} x - \frac{1}{2} (x - \tan^{-1} x)$$

$$= \frac{x^2}{2} \tan^{-1}x - \frac{x}{2} + \frac{1}{2} \tan^{-1}x + C$$

$$= \frac{1}{2} (x^2 + 1) \tan^{-1}x - \frac{x}{2} + C$$

$$= \left[\frac{1}{2} (x^2 + 1) \tan^{-1}x - \frac{x}{2} + C \right] \text{Ans}$$

Q-9 $\int_0^1 \frac{x}{x+1} dx$

$$= \int_0^1 \frac{x+1-1}{x+1} dx$$

$$= \int_0^1 \left(\frac{(x+1)}{x+1} - 1 \right) dx$$

$$= \int_0^1 \left(1 - \frac{1}{x+1} \right) dx$$

$$= \left[x - \log|x+1| \right]_0^1 \quad \int \frac{1}{x} dx = \log|x| + C$$

$$= [1 - \log(1+1)] - [0 - \log(0+1)]$$

$$= [1 - \log 2] - [-\log 1]$$

$$= 1 - \log 2 + \log 1 \quad (\because \log 1 = 0)$$

$$= [1 - \log 2] \quad \text{Ans}$$

$$\text{Q-10} \quad \int_0^1 \frac{(\log x)^3}{x} dx$$

$$= \int_0^1 (\log x)^3 \cdot \frac{1}{x} dx$$

$$= \left[\frac{(\log x)^4}{4} \right]_0^1 \quad \left(\because \frac{d}{dx} \log x = \frac{1}{x} \right)$$

$$= \frac{(\log 1)^4}{4} - \frac{(\log 0)^4}{4}$$

$$= \frac{(0)^4}{4} - \text{undefined}$$

This sum is undefined.

$$\text{Q-11} \quad \int_0^1 \frac{\tan^{-1} x}{1+x^2} dx$$

$$= \int (\tan^{-1} x)^1 \cdot \frac{1}{1+x^2} dx \quad \left[\int f(x)^n \cdot f'(x) dx = \int \frac{f(x)^{n+1}}{n+1} \right]$$

$$\left[\frac{(\tan^{-1} x)^2}{2} \right]_0^1$$

$$= \frac{1}{2} [\tan^{-1} 1]^2 - [\tan^{-1} 0]^2$$

Meditation is the best ode of worship.

$$= \frac{1}{2} [(\frac{\pi}{4})^2 - (0)^2]$$

$$= \frac{1}{2} (\frac{\pi}{4})^2$$

$$= \boxed{\frac{\pi^2}{8}} \quad \underline{\text{Ans}}$$

* Question set for 04 marks

$$1. \int \frac{1}{a^2 \cos^2 x + b^2 \sin^2 x} dx$$

$$= \int \frac{dx}{\cos^2 x (a^2 + b^2 \frac{\sin^2 x}{\cos^2 x})}, \quad (\because \frac{1}{\cos \theta} = \sec \theta,$$

$$= \int \frac{\sec^2 x \cdot dx}{a^2 + b^2 \tan^2 x}$$

$$= \frac{dy}{b^2 \left(\frac{a^2}{b^2} + \frac{y^2}{a^2} \right)}$$

$$= \frac{1}{b^2} \int \frac{du}{\left(\frac{a^2}{b^2} + u^2 \right)}$$

$$\int \frac{1}{x^2 + a^2} du = \frac{1}{a} \tan^{-1} \left(\frac{x}{a} \right) + C$$

પ્રભુને પ્રાર્થના એ શક્તિશાળી હૃથીયાર છે.

श्री रामनारायण डिवाईन मिशन

Dt.: / /

Pg.no.:

$$= \frac{1}{b^2} \times \frac{1}{a/b} \cdot \tan^{-1} \frac{u}{a/b} + C$$

$$= \frac{1}{b^2} \times \frac{b}{a} \tan^{-1} \left(\frac{bu}{a} \right) + C$$

$$= \left[\frac{1}{ab} \tan^{-1} \left(b \frac{tu}{a} \right) + C \right] \quad \underline{\text{Ans}}$$

Q3

$$\int \frac{\cos 3x}{\cos^2 x \cdot \sin^2 x} dx$$

Soln

$$= \int \frac{\cos^2 x - \sin^2 x}{\cos^2 x \cdot \sin^2 x} dx \quad (\because \cos 2x = \cos^2 x - \sin^2 x)$$

$$= \int \frac{\cos^2 x}{\cos^2 x \cdot \sin^2 x} - \frac{\sin^2 x}{\cos^2 x \cdot \sin^2 x} dx$$

$$= \int \frac{1}{\sin^2 x} - \frac{1}{\cos^2 x} dx$$

$$= \int \csc^2 x dx - \int \sec^2 x dx$$

$$= [-\cot x - \tan x + C] \quad \text{Ans}$$

મલ્યને પ્રાર્થના એ શક્તિશાળી હૃથીયાર છે.

$$\int \frac{1}{x(\log x - 1)(\log x - 2)} dx$$

$$\text{Let } \log x = t$$

$$\frac{1}{x} dx = dt$$

$$= \int \frac{dt}{(t-1)(t-2)}$$

$$= \int \frac{dt}{t^2 - 2(t)\left(\frac{3}{2}\right) + \left(\frac{3}{2}\right)^2 + 2 - \frac{9}{4}}$$

$$= \int \frac{dt}{\left(t - \frac{3}{2}\right)^2 - \left(\frac{1}{2}\right)^2}$$

$$= \boxed{\int \frac{1}{x^2 - a^2} dx = \frac{1}{2a} \log \left| \frac{x-a}{x+a} \right| + C}$$

$$= \frac{1}{2\left(\frac{1}{2}\right)} \cdot \log \left| \frac{t - \frac{3}{2} - \frac{1}{2}}{t - \frac{3}{2} + \frac{1}{2}} \right| + C$$

$$= \log \left| \frac{t + \left(-\frac{4}{2}\right)}{t + \left(\frac{-2}{2}\right)} \right| + C$$

$$= \log \left| \frac{t-2}{t-1} \right| + C$$

~~put the value of t in equation.~~
Meditation is the best ode of worship.

$$= \left[\log \left| \frac{\log x - 2}{\log x - 1} \right| + C \right] \quad \underline{\text{Ans}}$$

Q-5 $\int_0^{\pi/2} \frac{\sqrt{\sin x}}{\sqrt{\sin x} + \sqrt{\cos x}} dx$

$$= \int_0^{\pi/2} \frac{\sqrt{\sin x}}{\sqrt{\sin x} + \sqrt{\cos x}} dx \quad \textcircled{1}$$

$$= \int_0^{\pi/2} \frac{\sqrt{\sin(\pi/2 - x)}}{\sqrt{\sin(\pi/2 - x)} + \sqrt{\cos(\pi/2 - x)}} dx$$

$$= \int_0^{\pi/2} \frac{\sqrt{\cos x}}{\sqrt{\cos x} + \sqrt{\sin x}} dx \quad \textcircled{2}$$

Adding $\textcircled{1}$ and $\textcircled{2}$

$$2I = \int_0^{\pi/2} \frac{\sqrt{\sin x}}{\sqrt{\sin x} + \sqrt{\cos x}} + \frac{\sqrt{\cos x}}{\sqrt{\cos x} + \sqrt{\sin x}} dx$$

$$= \int_0^{\pi/2} \frac{\sqrt{\sin x} + \sqrt{\cos x}}{\sqrt{\sin x} + \sqrt{\cos x}} dx$$

$$\alpha I = \int_0^{\pi/2} 1 dx$$

$$\therefore 2I = \frac{\pi}{2}$$

$$2I = [x]_0^{\pi/2}$$

$$2I = \left[I = \frac{\pi}{4} \right] \text{ Ans}$$

$$\alpha I = \frac{\pi}{2} - 0$$

$$\text{Q-6} \quad \int_0^{\pi/4} \log(1 + \tan x) dx$$

$$= \int_0^{\pi/4} \log(1 + \tan(\frac{\pi}{4} - x)) dx$$

$$\left(\because \int_0^a f(x) dx = \int_0^a f(a-x) dx \right)$$

$$I = \int_0^{\pi/4} \log \left(\frac{1 + \tan \frac{\pi}{4} - \tan x}{1 + \tan \frac{\pi}{4} \cdot \tan x} \right) dx$$

$$\left(\because \tan(x-y) = \frac{\tan x - \tan y}{1 + \tan x \cdot \tan y} \right)$$

$$= \int_0^{\pi/4} \log \left(\frac{1 + \frac{1 - \tan x}{1 + 1 \cdot \tan x}}{1 + \tan x} \right) dx$$

$$= \int_0^{\pi/4} \log \left(\frac{1 + \tan x + 1 - \tan x}{1 + \tan x} \right) dx$$

Meditation is the best ode of worship.

$$= \int_0^{\pi/4} \log \left(\frac{2}{1 + \tan x} \right) dx$$

$$= \int_0^{\pi/4} \log 2 dx - \int_0^{\pi/4} (\underbrace{1 + \tan x}_{I = \text{given}}) dx$$

$$I + I = \int_0^{\pi/4} \log 2 \int_0^{\pi/4} 1 dx$$

$$2I = \log 2 (x) \Big|_0^{\pi/4}$$

$$= \log 2 \left(\frac{\pi}{4} - 0 \right)$$

$$= \log 2 \left(\frac{\pi}{4} \right)$$

$$2I = \frac{\pi}{4} \cdot (\log 2)$$

$$I = \frac{1}{2} \left(\frac{\pi}{4} \right) \log 2$$

$$\boxed{I = \frac{\pi}{8} \log 2} \quad \underline{\text{Ans}}$$

$$\text{Q-7} \quad \int_0^{\pi/2} \frac{\sqrt{\cot x}}{\sqrt{\tan x} + \sqrt{\cot x}} dx$$

$$\text{SOLM} \quad \int_0^{\pi/2} \frac{\sqrt{\cot x}}{\sqrt{\tan x} + \sqrt{\cot x}} dx \quad \text{Ans} \quad -\textcircled{1}$$

$$\int_0^{\pi/2} \frac{\sqrt{\cot(\pi/2 - x)}}{\sqrt{\tan(\pi/2 - x)} + \sqrt{\cot(\pi/2 - x)}} dx$$

$$= \int_0^{\pi/2} \frac{\sqrt{\tan x}}{\sqrt{\cot x} + \sqrt{\tan x}} dx \quad -\textcircled{2}$$

Adding $\textcircled{1}$ and $\textcircled{2}$

$$2I = \int_0^{\pi/2} \frac{\sqrt{\cot x} + \sqrt{\tan x}}{\sqrt{\tan x} + \sqrt{\cot x}} dx$$

$$\therefore 2I = \int_0^{\pi/2} \frac{\cancel{\sqrt{\cot x} + \sqrt{\tan x}}}{\cancel{\sqrt{\tan x} + \sqrt{\cot x}}} dx \quad 2I = \frac{\pi}{2} - 0$$

$$\therefore 2I = \int_0^{\pi/2} 1 dx$$

$$2I = \frac{\pi}{2}$$

$I = \frac{\pi}{4}$

Ans

$$\therefore 2I = [x]_0^{\pi/2}$$

Q8 Find the area of circle $x^2 + y^2 = a^2$

Soln centre = $(0, 0)$

radius = a

$$I = \int_a^b y dx$$

$$= \int_a^b F(x) dx$$

$$= \int_0^a \sqrt{a^2 - x^2} dx$$

$$\text{Formula: } \left[\int \sqrt{a^2 - x^2} dx = \frac{x}{2} \sqrt{a^2 - x^2} + \frac{a^2}{2} \sin^{-1} \frac{x}{a} \right]$$

$$= \left[\frac{x}{2} \sqrt{a^2 - x^2} + \frac{a^2}{2} \sin^{-1} \frac{x}{a} \right]_0^a$$

$$= \left[\frac{a}{2} \sqrt{a^2 - a^2} + \frac{a^2}{2} \sin^{-1} \frac{a}{a} \right] - \left[\frac{0}{2} \sqrt{a^2 - 0^2} + \frac{a^2}{2} \sin^{-1} \frac{0}{a} \right]$$

$$= \left[\frac{a}{2} (0) + \frac{a^2}{2} \sin^{-1}(1) \right] - \left[\frac{a^2}{2} (0) \right]$$

$$= \left[0 + \frac{a^2}{2} \cdot \frac{\pi}{2} \right] = 0$$

Area A = $\frac{1}{2} \pi a^2$

$$= \frac{a^2}{2} \cdot \frac{\pi}{2}$$

$$= \frac{a^2 \pi}{4}$$


$$= \frac{1}{4} a^2 \pi$$

$$\boxed{\text{Area} = \pi a^2}$$

Q9 Find the area enclosed by the parabola $y = x^2$ the x -axis and the line $x=2$ and $x=3$.

$$y = x^2$$

$x^2 = y$ which is a parabola



$$I = \int_a^b y \, dx$$

$$= \int_a^b f(x) \, dx$$

$$= \int_2^3 x^2 \, dx$$

$$= \left[\frac{x^3}{3} \right]_2^3$$

$$= \frac{1}{3} [x^3]_2^3$$

$$= \frac{1}{3} [(3)^3 - (2)^3]$$

$$= \frac{1}{3} [27 - 8]$$

$$= \frac{1}{3} [19]$$


$$I = \frac{19}{3}$$

$$\boxed{\text{Area} = \frac{19}{3}}$$

Ans

Q-1° Find the volume of sphere having radius a .

Soln



$$\text{circle } x^2 + y^2 = a^2$$

$$y^2 = a^2 - x^2$$

$$V = |I|$$

$$I = \pi \int_a^b y^2 dx \quad (\text{solution about } x = c \text{ is})$$

$$= \pi \int_a^b [F(x)]^2 dx$$

$$= \pi \int_{-a}^a [a^2 - x^2] dx$$

$a^2 - x^2$ is even function

$$= 2\pi \int_0^a [a^2 - x^2] dx$$

$$= 2\pi \left[a^2x - \frac{x^3}{3} \right]_0^a$$

$$= 2\pi \left[a^2 \cdot a - \frac{a^3}{3} \right] - 0$$

$$= 2\pi \left[\frac{3a^3 - a^3}{3} \right]$$

$$= 2\pi \left(\frac{2a^3}{3} \right)$$

$$\boxed{I = \frac{4\pi a^3}{3}}$$

volume of sphere =

$$V = |I|$$

$$= \boxed{\frac{4\pi a^3}{3}}$$

$$\boxed{\frac{4}{3} \pi a^3} \text{ Ans}$$